

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

B1

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

| | | |
|---|-----------|---|
| (51) International Patent Classification ⁶ : C12Q 1/68, 1/70, C12P 19/34, C07H 21/02, 21/04, G01N 33/53, C07K 15/28, 5/00 | A1 | (11) International Publication Number: WO 97/17470 (43) International Publication Date: 15 May 1997 (15.05.97) |
| (21) International Application Number: PCT/US96/17877 (22) International Filing Date: 8 November 1996 (08.11.96) (30) Priority Data: 08/555,394 9 November 1995 (09.11.95) US (71)(72) Applicant and Inventor: HOLLAND, James, F. [US/US]; 31 Mamaroneck Road, Scarsdale, NY 10583 (US). (72) Inventor: POGO, Beatriz, G., T.; 237 Nyac Avenue, Pelham, NY 10803 (US). (74) Agents: CLARK, Richard, S. et al.; Brumbaugh, Graves, Donohue & Raymond, 30 Rockefeller Plaza, New York, NY 10112 (US). | | (81) Designated States: AU, CA, JP, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> |
| (54) Title: DETECTION OF MAMMARY TUMOR VIRUS-LIKE SEQUENCES IN HUMAN BREAST CANCER (57) Abstract The present invention relates to materials and methods for diagnosing breast cancer in humans. It is based, at least in part, on the discovery that a substantial percentage of human breast cancer tissue samples contained nucleic acid sequences corresponding to a portion of the mouse mammary tumor virus <u>env</u> gene. In contrast, such sequences were absent in almost all other human tissues tested. | | |

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

| | | | | | |
|----|--------------------------|----|--|----|--------------------------|
| AM | Armenia | GB | United Kingdom | MW | Malawi |
| AT | Austria | GE | Georgia | MX | Mexico |
| AU | Australia | GN | Guinea | NE | Niger |
| BB | Barbados | GR | Greece | NL | Netherlands |
| BE | Belgium | HU | Hungary | NO | Norway |
| BF | Burkina Faso | IE | Ireland | NZ | New Zealand |
| BG | Bulgaria | IT | Italy | PL | Poland |
| BJ | Benin | JP | Japan | PT | Portugal |
| BR | Brazil | KE | Kenya | RO | Romania |
| BY | Belarus | KG | Kyrgyzstan | RU | Russian Federation |
| CA | Canada | KP | Democratic People's Republic of Korea | SD | Sudan |
| CF | Central African Republic | KR | Republic of Korea | SE | Sweden |
| CG | Congo | KZ | Kazakhstan | SG | Singapore |
| CH | Switzerland | LI | Liechtenstein | SI | Slovenia |
| CI | Côte d'Ivoire | LK | Sri Lanka | SK | Slovakia |
| CM | Cameroon | LR | Liberia | SN | Senegal |
| CN | China | LT | Lithuania | SZ | Swaziland |
| CS | Czechoslovakia | LU | Luxembourg | TD | Chad |
| CZ | Czech Republic | LV | Latvia | TG | Togo |
| DE | Germany | MC | Monaco | TJ | Tajikistan |
| DK | Denmark | MD | Republic of Moldova | TT | Trinidad and Tobago |
| EE | Estonia | MG | Madagascar | UA | Ukraine |
| ES | Spain | ML | Mali | UG | Uganda |
| FI | Finland | MN | Mongolia | US | United States of America |
| FR | France | MR | Mauritania | UZ | Uzbekistan |
| GA | Gabon | | | VN | Viet Nam |

Description

Detection Of Mammary Tumor Virus-Like Sequences In Human Breast Cancer

Cross-Reference to Related Application

This application is a continuation-in-part application of U.S. Serial No. 08/555,394, filed November 9, 1995.

Statement Regarding Federally Sponsored Research

- 5 This invention was made with funds from the U.S. government, which has certain rights in the invention.

Introduction

- 10 The present invention relates to materials and methods for diagnosing breast cancer in humans. It is based, at least in part, on the discovery that a substantial percentage of human breast cancer tissue samples contained nucleic acid sequences corresponding to a portion of the mouse mammary tumor virus env gene. In contrast, such sequences were absent in almost all other human tissues tested.
- 15

Background of the Invention

- 20 A large body of information has accumulated about the molecular biology of MMTV (reviewed in Slagle, B.L. et al., 1987, in "Cellular and Molecular Biology of Mammary Cancer", Kidwell et al., eds., Plenum Press, NY. pp 275-306). Mouse mammary tumor virus (MMTV) is associated with a high incidence of breast cancer in certain strains of mice (over 90% among females), and has been regarded as a potential model for human disease.
- 25

The MMTV virus does not carry a transforming oncogene, but rather acts as an insertional mutagen with several proviral insertion loci designated int-1

or wnt-1 (Nusse R. et al., 1982, Cell 31:99-109) int-2
(Peters, G. et al., 1983, Cell 33:369-377) int-3
(Gallahan, D. et al., 1987, J. Virol. 61:218-220) int-4
(Roelink, H. et al., 1990, Proc. Natl. acad. Sci. USA
5 87:4519-4523) and int-5 (Morris, V.L., et al. 1991,
Oncogene Research 6:53-63), which encode for growth
factors or other related proteins. These genes are not
expressed in normal mammary tissue but become activated
after integration of MMTV provirus into the adjacent
10 chromosomal DNA.

The human homolog of the int-2 locus has been
located on chromosome 11 (Casey, G. et al., 1986,
Mol. Cell Biol. 6:502-510) and has been found amplified
(in 15% of the breast cancers) and also expressed
15 (Lidereau, R. et al., 1988, Oncogene Res 2:285-291;
Zhou, D.J. et al., 1988, Oncogene 2:279-282; Liscia,
D.S. et al., 1989, Oncogene 4:1219-1224; Meyers, S.L.
et al., 1990, Cancer Res 50:5911-5918). It may be
significant that in tumors from Parsi women, who have a
20 high incidence of breast tumors, the int-2 locus is
amplified in 50% of the cases (Barnabas-Sohi, N. et
al., 1993, Breast Dis. 6:13-26). The amplification of
int-2 and other genes in 11q13 is indicative of poor
prognosis (Schuwring, E. et al., 1992, Cancer Research
25 52:5229-5234; Champeme, M-H, et al., 1995, Genes,
Chromosomes and Cancer 12:128-133). Both mouse and
human int-2 have been sequenced (Moore, R. et al.,
1986, EMBO J 5:919-924). The gene encodes a protein of
about 27 kilodaltons (KD) which shows homology to both
30 basic and acidic fibroblast growth factors (Dickson, C.
et al. 1987, Nature (London) 326:833).

However, efforts to demonstrate the presence of
viruses in human breast cancer through search for viral
particles, immunological cross-reactivity, or sequence
35 homology have yielded contradictory results. Detect-
able MMTV env gene-related antigenic reactivity
has been found in tissue sections of breast cancer

(Mesa-Tejada et al., 1978, Proc. Natl. Acad. Sci. USA 75:1529-1533; Levine, P. et al., 1980, Proc. Am. Assoc. Cancer Res. 21:170; Lloyd, R. et al., 1983, Cancer 51:654-661), breast cancer cells in culture (Litvinov, S.V. and Golovkina, T.V., 1989, Acta Virologica 33:137-142), human milk (Zotter S. et al., 1980, Eur. J. Cancer 16:455-467) in sera of patients (Day, N.K. et al., 1981, Proc. Natl. Acad. Sci. USA 78:2483-2487), in cyst fluid (Witkin, S.S. et al., 1981, J. Clin. Invest. 67:216-222) and in particles produced by a human breast carcinoma cell line (Keydar, I. et al., 1984, Proc. Natl. Acad. Sci. USA 81:4188-4192). Sequence homology to MMTV has been found in human DNA under low stringency conditions of hybridization (Callahan, R. et al., 1982, Proc. Natl. Acad. Sci. USA 79:5503-5507) and RNA related to MMTV has been detected in human breast cancer cells (Axel, R. et al., 1972, Nature 235:32-36). The presence of MMTV related sequences in lymphocytes from patients with breast cancer has been reported (Crepin, M. et al., 1984, Biochem. Biophys. Res. Comm. 118:324-331), as well as detection of reverse transcriptase (RT) activity in their monocytes (Al-Sumidaie, A.M. et al., 1988, Lancet 1:5-8). May and Westley (May and Westley, 1989, Cancer Research 49:3879-3883) have reported the presence of MMTV-like sequences arranged as tandem repeats only in DNA from breast cancer cells.

These results have been difficult to interpret, and theories linking MMTV or a related virus with human breast cancer have fallen out of favor, in view of the relatively recent discovery of human endogenous retroviral sequences ("HERs"; Westley, B. et al., 1986, J. Virol. 60:743-749; Ono, M. et al., 1986, J. Virol. 60:589-598; Faff, O. et al., 1992, J. Gen. Virology 73:1087-1097). Data which could be interpreted to demonstrate the presence of MMTV-related sequences could be more readily explained by endogenous human

retroviral sequences. Adding further confusion to the picture, env-gene related antigenicity has been detected in epitopes of human proteins (Hareuveni, M. et al., 1990, Int. J. Cancer 46:1134-1135).

5 Brief Summary of the Invention

 The present invention relates to methods for diagnosing breast cancer in humans in which the presence of mouse mammary tumor virus env gene-like sequences bears a positive correlation to the existence
10 of malignant breast disease. It is based, at least in part, on the discovery that 38 to 40 percent of human breast cancer tissue samples tested contained gene sequences homologous to the mouse mammary tumor virus env gene that are substantially absent from other human
15 tumors and tissues. The invention also relates to methods for diagnosing breast cancer in humans in which the presence of retrovirus proviral fragments substantially homologous to the env gene and/or 3' LTR sequence of MMTV are detected. The molecular probes
20 used in these experiments were designed to avoid cross-hybridization with endogenous human retroviral sequences. The present invention further provides for compositions of molecular probes which may be utilized in such diagnostic methods.

25 Brief Description of the Figures

FIGURE 1: Amplification of 660 bp of MMTV-like env gene. DNA was extracted from frozen tissues. PCR was performed using primers 1 and 3. A: 2% agarose gel electrophoresis. B: Southern blot hybridization
30 using 5'³²P-end-labeled probe 2. Lanes 1 and 3: breast cancer; lanes 2 and 4: normal breast; lane 5: control reaction (no DNA); lane E: MMTV env gene. M: molecular weight marker. Arrow indicates 510 bp band.

FIGURE 2: Nested PCR. A: 2% agarose gel electrophoresis. 1: Amplification of 686 bp of MMTV-like env
35

gene sequences using primers 1 and 4 and the product of reaction A 1 as template. 2: Amplification of 250 bp of MMTV-like env gene sequences using primers 2 and 3. B, 1 and 2: Southern blot hybridization of the amplified products using probe 5'-³²P end-labeled probe 2a.

FIGURE 3: Amplification of 250 bp of MMTV-like env gene. DNA was extracted from paraffin-embedded tissue sections. PCR was performed using primers 2 and 3. A: 2% agarose gel electrophoresis. B: Southern blot hybridization using 5'-³²P-labeled probe 2a. Lane 1: normal breast; lanes 2 to 5: breast cancer; lane E: MMTV env gene. M: molecular weight marker. Arrow indicates 298 bp band.

FIGURE 4: Nucleotide sequence of the cloned MMTV env gene-like sequences as compared to the env sequences of the GR and BR6 strains of MMTV using the GCG program. *:potential glycosylation site, |:mismatch to MMTV.

FIGURE 5: Southern blot hybridization of genomic DNA. DNA was extracted from frozen tissues or cell lines, digested with EcoRI and transferred to nitrocellulose paper. Hybridization with ³²P-labeled clone 166. DNA from A, B, and G: env gene positive breast cancer; C and D: env negative breast cancer; E and F: normal breast; H: MCF-7 cells. M: molecular weight marker, Arrow indicates 9kb band.

FIGURE 6: Southern blot hybridization of genomic DNA. Experimental conditions as in Fig. 5. DNA from A and B: env negative breast cancer; C and D: env positive breast cancer; E: molecular weight marker (non-labelled); F. to H: normal breast. Arrow indicates position of 9 kb marker.

FIGURE 7: Map of MMTV.

FIGURE 8: Comparison of the nucleic acid sequence of mouse mammary tumor env gene ("MMTENV"), showing residues 976-1640, with the nucleic acid sequence of a

representative 660 bp sequence obtained by PCR reaction of DNA from human breast cancer tissue ("MS1627").

FIGURE 9: Sequence of an about 2.6 kb MMTV-like fragment detected in a human breast carcinoma.

5 Detailed Description of the Invention

The present invention relates to methods and compositions for diagnosing breast cancer in humans.

The present invention provides for compositions comprising an isolated and purified nucleic acid

10 molecule which (i) hybridizes to a gene of mouse mammary tumor virus; (ii) is present in at least 20 percent of DNA samples prepared from breast cancer tissue of different human subjects; and (iii) is present in less than 5 percent of DNA samples prepared

15 from tissues other than breast cancer tissue from different human subjects. A "gene of mouse mammary tumor virus" includes, but is not limited to, the gag, pol, and env genes and the 5' LTR and 3' LTR sequences of MMTV. In preferred embodiments of the invention,

20 the mouse mammary tumor virus (hereafter "MMTV") gene is the env gene and/or the 3' LTR sequence. The term "hybridize" is used to refer to routine DNA-DNA or DNA-RNA hybridization techniques under what would be regarded, by the skilled artisan, as stringent

25 hybridization conditions. The phrase "is present" indicates that a native form of the molecule, in an unpurified state (for example, as part of chromosomal DNA), may be detected by a standard laboratory technique, such as Southern blot or polymerase chain

30 reaction (PCR). To be "present", the molecule may be detectable by one technique but not others. To be present in "less than 5 percent of DNA samples prepared from tissues other than breast cancer tissue from different human subjects", all non-breast cancer tissue

35 samples are considered together, but the total number of samples must be large enough to give the 5 percent

value statistical significance that would be reasonable to the skilled artisan.

In order to identify such a nucleic acid molecule, the sequence of MMTV may be compared, using a computer database, to known human DNA sequences, and portions of MMTV which are less than or equal to 25 percent homologous to a human sequence may be selected for further study. The term "homologous", as used herein, refers to the presence of identical residues; for example, a first sequence is considered 25 percent homologous to a second sequence if it shares 25 percent of the residues of the first sequence. Since there is relatively greater likelihood that MMTV may bear similarity to human retroviral-like sequences, it may be preferable to evaluate whether a particular MMTV nucleic acid sequence is homologous to such sequences, for example, as endogenous human retrovirus sequences. A prototype of such viruses is HERV-K10 (Ono, M. et al., 1986, J. Virol. 60:589-598).

Once an MMTV gene sequence which is less than or equal to 25 percent homologous to a human DNA sequence, such as a human endogenous retroviral sequence, is identified, the presence of nucleic acid molecules having the MMTV gene sequence in human breast cancer tissues and other tissues may be evaluated. Such evaluations may be performed either by Southern blot techniques, or, preferably, by polymerase chain reaction (PCR) techniques, which are more sensitive. In such a way, MMTV gene sequences which (i) hybridize to at least 20 percent of DNA samples prepared from breast cancer tissue of different human subjects and (ii) hybridize to less than 5 percent of DNA samples prepared from human tissues other than breast cancer tissues may be identified. A nucleic acid molecule having a MMTV gene sequence which satisfies these requirements may then be used in diagnostic methods which detect the presence of such sequence in human

breast tissue by standard techniques, including PCR techniques which assay for the presence of the molecule, but also, where appropriate, Southern blot, Northern blot, or Western blot techniques, to name but a few.

In preferred embodiments, the present invention relates to a portion of MMTV localized between MMTV env gene sequences 976 and 1640 (Majors, I.E. and Varmus, H.E., 1983, J. Virol. 47:495-504; see Fig. 7). This about 660 bp sequence (hereafter, "the 660 bp sequence") has been found to exhibit low (16 percent) homology to the prototype human endogenous retrovirus HERV-K10, using the IBI/Pustell Sequence Analysis Program, and has also been shown to be present in 121 (38.5%) of 314 unselected breast cancer tissue samples, in cultured breast cancer cells, in 2 of 29 breast fibroadenomas (6.9%) and in 2 of 107 breast specimens from reduction mammoplasties (1.8%). The sequence was not found in normal tissues including breast, lymphocytes from breast cancer patients nor in other human cancers or cell lines (see example section, infra). Similarly, an about 250 bp sequence (hereafter "the 250 bp sequence"), between positions 1388 and 1640 in the env gene, and therefore falling within the 660 bp sequence, was detected in 60 (39.7%) of 151 breast cancer, and in one of 27 normal breast samples assayed from paraffin-embedded sections. Cloning and sequencing of the 660 bp and 250 bp sequences demonstrated that they are 95-99% homologous to MMTV env gene, but not to the known human endogenous retroviruses ("HERs") nor to other viral or human genes (<18%).

In another preferred embodiment, the present invention relates to a nucleic acid molecule which corresponds to a retroviral genomic fragment which has substantial homology to 3' LTR and/or env gene of the MMTV genome, and is found in a substantial percentage of breast cancer samples. By substantial percentage is

meant at least 20% of tested breast cancer samples.

Such a sequence is preferably comprised of the 3' LTR region and all or part of the env gene, although it may include more sequences of a retroviral genome. Most preferably, the sequence is at least comprised of an about 2.6 kb fragment which comprises the 1,228 base pair (bp) sequence of the 3' LTR sequence and 1,336 bp of the env gene sequence of MMTV (Fig. 9) (SEQ ID NO:20). When compared with the two strains of MMTV C3H and BR6, the sequence homology was 90.8% and 90.7%, respectively. When compared with the endogenous retroviral sequences (HUMERKA), sequence homology was only 58% in 36 bp and 71% in 74 bp.

Retrovirus proviral sequences can be detected by PCR technology using primers derived from the MMTV genome. Such primers include primer 5L, containing the nucleotides 7376-7395 of the MMTV BR6 genome (5'-3': CCAGATCGCCTTTAAGAAGG) (SEQ ID NO:11) and primer LTR3, containing nucleotides 9918-9927 of the MMTV BR6 genome (5'-3': CGAACAGACACAAAGCGACG) (SEQ ID NO:19). Other primers which correspond to or are homologous to MMTV sequences can be used as primers. Nucleotide fragments which correspond to or are homologous to the retroviral sequences isolated from the breast cancer samples can also be used to amplify additional retroviral fragments from the samples. Long PCR techniques can be used to amplify longer stretches of a proviral sequence.

The present invention provides for compositions comprising an isolated and purified nucleic acid molecule which hybridizes to the about 2.6 kb retroviral fragment shown in Fig. 9 under stringent conditions or is at least 90 percent homologous to said fragment using the MacVector homology determining program which may be used to diagnose breast cancer in a subject, using methods which include PCR and Southern blot methods.

Nucleic acids having the 660 bp sequence, the 250 bp sequence, or all or part of the about 2.6 kb sequence, may therefore be used, according to the invention, to diagnose breast cancer in a subject, using methods which include PCR and Southern blot methods. Where PCR methods are used, primers such as those listed in Table 1, below, may be utilized.

The present invention provides for compositions comprising essentially purified and isolated nucleic acid having the 660 bp sequence or the 250 bp sequence or an at least five bp, and preferably greater than or equal to ten bp, subsequence thereof. In order to maintain the desired specificity, such nucleic acid molecules may preferably contain sequence falling within the 660 bp sequence, but preferably do not contain sequences from other portions of the MMTV genome, which may, undesirably, hybridize to human sequences which are not breast cancer specific, such as HERs. Accordingly, the present invention provides for compositions wherein the isolated and purified nucleic acid molecule comprises at least a portion having a nucleic acid sequence which hybridizes to a region of the mouse mammary tumor virus env gene between residues 976 and 1640, or between residues 1388 and 1640, and wherein the isolated and purified nucleic acid molecule does not hybridize to any other region of the MMTV genome.

The 660 bp sequence, in various embodiments, may have a number of nucleotide sequences. For example, in one embodiment, the 660 bp sequence may have a sequence as set forth in Fig. 8 and designated "MMTENV-like sequence" (SEQ ID NO:17), which depicts the MMTV env sequence between residues 976 and 1640. In a second series of embodiments, the 660 bp sequence may have a sequence as set forth in Fig. 8 and designated "MS1627" (SEQ ID NO:18), which depicts a predominant sequence for the 660 bp sequence as it has been defined by

sequencing analysis of the products of PCR reactions using DNA from human breast cancer tissues. In still further embodiments, the 660 bp sequence may have various other nucleotide sequences obtained by
5 sequencing the results of PCR reactions to detect the presence of 660 bp sequence in human breast cancer tissues.

In related embodiments, the present invention provides for compositions comprising PCR primers
10 that may be used to detect the presence of the forementioned molecules or other MMTV-like sequences. For example, the compositions may comprise one or more of the following primer molecules (5' - 3'):
CCTCACTGCCAGATC (SEQ ID NO:1); GGGAATTCCTCACTGCCAGATC
15 (SEQ ID NO:2); CCTCACTGCCAGATCGCCT (SEQ ID NO:3);
TACATCTGCCTGTGTTAC (SEQ ID NO:4); CCTACATCTGCCTGTGTTAC
(SEQ ID NO:5); CCGCCATACGTGCTG (SEQ ID NO:6);
ATCTGTGGCATACT (SEQ ID NO:7); GGGAATTCATCTGTGGCATACT
(SEQ ID NO:8); ATCTGTGGCATACTTAAAGG (SEQ ID NO:9);
20 GAATCGCTTGGCTCG (SEQ ID NO:10); CCAGATCGCCTTTAAGAAGG
(SEQ ID NO:11); TACAGGTAGCAGCACGTATG (SEQ ID NO:12);
CGAACAGACACAAACACACG (SEQ ID NO:19).

The use of such compositions and molecules in PCR and Southern blot techniques is illustrated in the non-
25 limiting examples set forth below. The correlation between the presence of the MMTV-related nucleic acid molecules described above and breast cancer allows such molecules and compositions to be utilized in the diagnosis of breast cancer. Accordingly, the present
30 invention provides for a method of diagnosing breast cancer, wherein the detection of such nucleic acid molecules bears a positive correlation to the existence of breast cancer in a human. The results of such
evaluation, together with additional clinical symptoms,
35 signs, and laboratory test values, may be used to formulate the complete diagnosis of the patient.

In further related embodiments, the present invention provides for an essentially purified peptide encoded by a nucleic acid molecule which (i) hybridizes to a gene of MMTV; (ii) is present in at least
5 20 percent of DNA samples prepared from breast cancer tissue of different human subjects; and (iii) is present in less than 5 percent of DNA samples prepared from tissues other than breast cancer tissue from different human subjects. In preferred embodiments, the
10 MMTV gene is the env gene.

Such peptides may be used in the diagnosis of breast cancer. Accordingly, the present invention provides for a method of diagnosing breast cancer in a human subject, comprising detecting the presence of
15 a peptide encoded by a nucleic acid molecule which (i) hybridizes to the env gene of a mouse mammary tumor virus; (ii) is present in at least 20 percent of DNA samples prepared from breast cancer tissue of different human subjects; and (iii) is present in less than
20 5 percent of DNA samples prepared from tissues other than breast cancer tissue from different human subjects.

The present invention also provides for antibodies (including monoclonal and polyclonal) antibodies which
25 specifically bind to such peptides. Such antibodies may be used in methods of diagnosing breast cancer, for example, but not by way of limitation, by Western blot, immunofluorescent techniques, and so forth.

In nonlimiting embodiments of the invention, the
30 skilled artisan may evaluate MMTV-like nucleic acid molecules for regions which would be considered likely to encode immunogenic peptides (using, for example, hydropathy plots). Such peptides may then be sequenced and used to produce antibodies that may be employed in
35 diagnostic methods as set forth above.

For example, certain peptides encoded by portions of the 660 bp sequence have been synthesized. These

peptides, which have the sequences LKRPGFQEHMI (SEQ ID NO:13) and GLPHLIDIEKRG (SEQ ID NO:14), have been used to produce antibodies in rabbits, and the resulting antisera have successfully identified breast cancer cells positive for MMTV env-like sequences by PCR assay. Other peptides encoded by 660 bp sequence which may be useful according to the invention include TNCLDSSAYDTA (SEQ ID NO:15) and DIGDEPWFD (SEQ ID NO:16).

10 6. Example: The Detection of Mouse Mammary Tumor Virus Env Gene-Like Sequences in Human Breast Cancer Cells and Tissues

 6.1. Materials and Methods

 DNA from breast cancer tissue and other human cancer tissues, human placentas, normal human tissues including breast, and from several human cell lines (including eight breast cancer cell lines), and two normal breast cell lines was extracted following the procedure of Delli Bovi et al. (1986, Cancer Res. 46:6333-6338). The DNA was resuspended in a solution containing 0.05 M Tris HCl buffer, pH 7.8, and 0.1 mM EDTA, and the amount of DNA recovered was determined by microfluorometry using Hoechst 33258 dye (Cesarone, C. et al., 1979, Anal Biochem 100:188-197). Plasmids containing the cloned genes of MMTV were obtained from the ATCC, propagated in Escherichia coli cultures and purified using anion-exchange minicolumns (Qiagen) or by precipitation with polyethylene glycol (Sambrook J., et al., 1989, in "Molecular Cloning/A Laboratory Manual", Cold Spring Harbor). Oligonucleotide primers were synthesized at the core facilities of the Brookdale Molecular Biology Center at Mount Sinai School of Medicine.

 Polymerase chain reaction (PCR) was performed using Taq polymerase following the conditions recommended by the manufacturer (Perkin Elmer Cetus)

with regard to buffer, Mg^{2+} and nucleotide concentrations. Thermocycling was performed in a DNA cycler by denaturation at 94° C for 3 min. followed by either 35 or 50 cycles of 94° C for 1.5 min., 50° C for 2 min. and 72° C for 3 min. The ability of the PCR to amplify the selected regions of the MMTV env gene was tested by using as positive templates the cloned MMTV env gene and the genomic DNA of the MCF-7 cell line, since it was shown to express gp52 immunological determinants (Yang, N.S., et al., 1975, J. Natl. Cancer Inst. 61:1205-1208). Optimal Mg^{2+} , primer concentrations and requirements for the different cycling temperatures were determined with these templates. The master mix as recommended by the manufacturer was used. To detect possible contamination of the master mix components, a reaction without template was routinely tested. γ DNA and control primers provided by the manufacturer were used as control for polymerase activity. As an internal control, amplification of a 120 bp sequence estrogen receptor gene was assayed using primers designed and generously provided by Dr. Beth Schachter, (Mount Sinai School of Medicine, N.Y.). In addition, primers for actin 5 gene amplification were also used.

The product of the PCR was analyzed by electrophoresis in a 2% agarose gel. A 1 kb DNA ladder (Gibco BRL) was used to identify the size of the PCR product. To determine if the amplified sequences of the middle region of the 660 bp faithfully reproduced the sequences of the env gene of MMTV, an 18-mer sequence within the env gene was used as a probe for the 660 bp amplified sequence. The 18-mer probe was 5' end-labeled with ^{32}P -ATP using T4 polynucleotide kinase and purified by the NENSORB nucleic acid purification cartridge (NEN). Southern blot hybridization was performed using the conditions described by (Saiki et al., 1985, Science 230:1350-1354).

The product of the PCR (660 bp or 250 bp) was cloned directly from the reaction mixture into the TA cloning vector (Invitrogen) using the TA cloning kit and following the conditions recommended by the supplier. Direct cloning of the fragment isolated from the gel, was also performed. Plasmid DNA was purified by CsCl density gradient centrifugation or by precipitation with polyethylene glycol (Sambrook et al., 1989, in "Molecular Cloning/A Laboratory Manual", Cold Spring Harbor), restricted with HindIII and EcoRI, electrophoresed in 2% agarose gels and transferred to nitrocellulose filters. Southern blot hybridization was carried out using a 5'-terminal labeled internal probe as described above. Cloning procedures were performed in laboratories totally separate from those where PCR was carried out. Automated DNA sequencing (using Applied Technology Sequencer Model 373A) was performed in the Brookdale Molecular Biology Center. Sequence homology was determined using the IBI MacVector GenBank and GCG Programs.

To prevent contamination of the samples, processing of human tissues was performed in a laminar flow hood. DNA extractions were done in a chemical hood located in a different room from that where PCR was performed. PCR assays were assembled in a biological hood provided with ultraviolet light. Aerosol resistant tips and dedicated positive-displacement pipettes were used throughout. All equipment used for PCR (microcentrifuge, electrophoresis apparatus, pipettors) was cleaned each time with 10% sodium hypochlorite to assure DNA decontamination (Prince and Andrus, 1992, Biotechniques 12:358-36). After the initial experiments were performed, the plasmid containing the MMTV env gene was frozen and never used again, to avoid contamination. However, to detect plasmid contamination from our own env gene clones,

primers were designed to amplify plasmid sequences. All the authentic MMTV env positive samples were then tested and found negative for plasmid contamination.

Southern blotting and hybridization were performed
5 as described (Southern, E.M., 1975, J. Mol. Biol. 98:503-517), using the 660 bp cloned sequences labeled by the random primer procedure (Feinberg, A.P., et al., 1983, Anal. Biochem. 132:6-13). Prehybridization and hybridization were performed in a solution containing
10 6 x SSPE, 5% Denhardt's, 0.5% SDS, 50% formamide, 100 µg/ml denaturated salmon testis DNA, incubated for 18 hrs at 42°C, followed by washings with 2 x SSC and 0.5% SDS at room temperature and at 37°C and finally in 0.1 x SSC with 0.5% SDS at 68°C for 30 min (Sambrook
15 et al., 1989, in "Molecular Cloning/A Laboratory Manual", Cold Spring Harbor). For paraffin-embedded tissue sections the conditions described by Wright and Manos (1990, in "PCR Protocols", Innis et al., eds., Academic Press, pp. 153-158) were followed using
20 primers designed to detect a 250 bp sequence.

6.2. Results

6.2.1. Selection of Specific MMTV Env Gene Sequences

A computer search for MMTV env gene homologous sequences was first performed, since sequence homology
25 between the human endogenous retroviral sequences and MMTV had been described. The prototype of this group of human endogenous retroviruses is HERV-K10 (Ono, M. et al., 1986, J. Virol. 60:589-598). The sequences of the env gene of MMTV (Majors, I.E. and Varmus, H.E.,
30 1983, J Virol 47:495-504) were aligned with sequences of the env gene of the human endogenous retrovirus HERV-K10 (Ono, M. et al., 1986, J. Virol. 60:589-598), using the IBI/Pustell Sequence Analysis Program. A region of 660 bp of low homology (16%) was localized
35 between MMTV env gene sequences 976 and 1640 (Majors, I.E. and Varmus, H.E., 1983, J Virol 47:495-504). This

internal domain of the outer membrane of the env gene has only one glycosylation site and is highly conserved between strains. Two primers comprising 15 bp sequences at positions 976-990 (primer 1) and 1626-1640 (primer 3) were first synthesized. Later longer primers were synthesized (1N and 3N). An 18-mer sequence in the middle of the 660 bp MMTV env region (1388-1405) (primer 2) was used as a probe to identify the 660 bp sequence. A second oligomer probe was synthesized comprising the sequence 1554 to 1568 (primer 2a) to be used for hybridization when a sequence of around 250 bp (between positions 1388 and 1640) was amplified. For nested PCR reactions (Mullis, K.B. and Faloona, F.A., 1987, Meth Enzymol 155:335-350), another primer comprising sequences 1647 to 1661 (primer 4) was synthesized to be used with primer 1 in the first reaction and primers 2 and 3 in the second. Modified primers with GC clamps and extra sequences were also synthesized and used in the PCR (primers 1a and 3a). Another set of primers comprising sequences 974 to 1003 (5L) and 1558 to 1577 (3L) were subsequently developed because their T_m's matched and provided better amplification than the original primers. The sequences are represented in Table 1. All of them were productive in amplification reactions.

Table 1. Primer and probe sequences and location in mouse mammary tumor virus env gene

| | Designation | Sequence (5'-3') | Location |
|----|-------------|------------------------|-----------|
| 5 | 1 | CCTCACTGCCAGATC | 976-990 |
| | 1a | GGGAATTCCTCACTGCCAGATC | 976-990 |
| | 1N | CCTCACTGCCAGATCGCCT | 976-993 |
| | 2 | TACATCTGCCTGTGTTAC | 1388-1405 |
| 10 | 2N | CCTACATCTGCCTGTGTTAC | 1386-1405 |
| | 2a | CCGCCATACGTGCTG | 1554-1568 |
| | 3 | ATCTGTGGCATACT | 1640-1626 |
| | 3a | GGGAATTCATCTGTGGCATACT | 1640-1626 |
| | 3N | ATCTGTGGCATACTAAAGG | 1640-1621 |
| 15 | 4 | GAATCGCTTGGCTCG | 1661-1647 |
| | 5L | CCAGATCGCCTTTAAGAAGG | 984-1003 |
| | 3L | TACAGGTAGCAGCACGTATG | 1558-1577 |

6.2.2. Detection of MMTV-Like Env Gene Sequences in Human Breast Tumor DNA

20 PCR was performed on DNA extracted from breast cancer tissues, normal breast tissues and from the plasmid containing the env gene of MMTV, using primers 1 and 3. Photographs of the ethidium bromide stained gels of the PCR product reveal the presence of an

25 approximately 660 bp sequence in some of the tumors, (Fig. 1A, lanes 1 and 3) but not in the normal tissue samples (Fig. 1A, lanes 2 and 4). As a positive control the MMTV env gene was also amplified (Fig. 1A, lane E). Similar results were obtained with modified

30 primers 1a, 3a, 3L and 5L. Southern blot hybridization of the gel with ³²P-labeled 18-mer oligonucleotide (primer 2) indicated that this internal sequence was present in the amplified material (Fig. 1B) and that the bands in the gel were not artifactual.

35 Our initial effort was to analyze a representative sample of breast cancer specimens as well as normal

tissues and other tumors. To date 343 breast tumors have been processed, DNA extracted and PCR performed. Of these 343 tumors, 314 were carcinomas and 29 were fibroadenomas. Amplification of sequences of 660 bp was observed in 121 of the carcinomas (38.5%) and in 2 of the 29 fibroadenomas (6.9%). These sequences were confirmed to be MMTV env gene-like sequences by hybridization with the labeled specific probe containing the internal sequences. These sequences were not detected in the DNAs extracted from 20 normal organs, 23 cancers from other organs and 26 samples of blood lymphocytes including 7 from breast cancer patients whose breast specimens were positive. From 107 samples of normal breast obtained from reduction mammoplasties, 2 were positive (1.8%). In addition to DNA from lymphocytes from seven positive patients, DNA from their normal breast tissue of the operated breast was tested in 4 cases. All were negative (Table 2). Finally, DNA of the MCF-7, and ED (a cell line developed in our laboratory from the pleural effusion of a patient with an env -positive breast tumor) breast cancer cell lines were shown to contain the 660 bp MMTV env gene-like sequences (Table 3), while four other breast cancer cell lines were positive only for the 250 bp sequence (T47-D, BT-474, BT-20 and MDA-MB-231).

Table 2. Detection of MMTV env gene-like sequences in human DNA extracted from fresh or frozen tissues

| 5 | <hr/> | | | |
|-------|---|--------|--------------------------------|------------|
| | Sample | Number | MMTV <u>env</u> gene sequences | % Positive |
| <hr/> | | | | |
| | Breast Carcinomas | 314 | 121 | 38.5% |
| | Breast Fibroadenomas | 29 | 2 | 6.9% |
| 10 | Normal Breasts | 107 | 2 | 1.8% |
| | *Normal Breasts | 4 | negative | |
| | Tumors other than breast | 23 | negative | |
| | Normal tissues | 20 | negative | |
| 15 | Lymphocytes | 26 | negative | |
| | **Lymphocytes | 7 | negative | |
| <hr/> | | | | |
| | * Histologically normal tissue from same breast as positive cancer. | | | |
| 20 | ** Lymphocytes from breast cancer patients who were positive for MMTV <u>env</u> gene sequences in the tumor. | | | |

Table 3. Detection of MMTV env gene-like sequences in DNA from human cell lines in culture

| | Human Cell Lines | | MMTV <u>env</u> gene sequence |
|----|------------------|--------------------------|-------------------------------|
| 5 | MC-7 | (breast carcinoma) | positive |
| | T47-D | " " | negative |
| | BT-20 | " " | negative |
| | MDA-MB-231 | " " | negative |
| | ZR-75-1 | " " | negative |
| 10 | SK-BR 3 | " " | negative |
| | BT474 | " " | negative |
| | ED | " " | positive |
| | MCF-10 | (normal breast) | negative |
| | HB-447 | " " | negative |
| 15 | HL-60 | (promyelocytic leukemia) | negative |
| | K562 | (erythroleukemia) | negative |
| | Jurkat | (T cell leukemia) | negative |
| | Hep 6-2 | (hepatoma) | negative |

The nested polymerase reaction was used in several instances to increase sensitivity and specificity, thus reducing the probability of false positives. In Fig. 2, results of a representative nested reaction are shown using primers 1 and 4 in the first reaction (Fig. 2A) and 2 and 3 for the 2nd reaction. The specificity of the reaction can be seen in the 2nd amplification (Fig. 2B).

To study a large number of samples and to be able to perform archival studies, PCR of paraffin-embedded tissue sections was also carried out. Primers 2 and 3 were used to amplify a 250 bp sequence within the 660 bp stretch when DNA was extracted from paraffin-embedded tissue sections since larger size sequences are difficult to amplify after fixation. Tumor DNA was amplified (Fig. 3A, lanes 2-5) whereas normal breast DNA was not (Fig. 3A, lane 1). The identification of

this 250 bp sequence with the MMTV-like env gene was confirmed by hybridization with an internal probe (primer 2a) as shown in Fig. 3B. Using this procedure we have analyzed 151 breast cancer samples and found
5 that 60 (39.7%) possess the 250 bp sequence. Of the 27 normal breast samples obtained from reduction mammoplasties assayed by this procedure, one was positive (3.7%). These results, in conjunction with those obtained from lymphocytes and from normal breast
10 tissue of patients whose breast cancer was PCR positive, indicate that MMTV-like sequences are present in a significant number of human breast cancer DNA which cannot be explained by DNA polymorphism.

15 6.2.3. Cloning and Sequencing of the
MMTV-Like Env Gene Sequences

To find out whether there was homology to MMTV env gene throughout the whole 660 bp stretch, the product of the PCR from 8 different tumors was cloned and sequenced. In Fig. 4 the sequence of different clones
20 comprising around 600 bp are represented, as aligned to the MMTV env gene sequence of the GR and BR6 strains (Redmon, S. and Dickson, C., 1983, EMBO J. 2:125-131). This domain of the env gene in the GR strain is 100% homologous to the C₃H strain and 98% to the BR6 strain
25 (Majors, I.E. and Varmus, H.E., 1983, J. Virol. 47:495-504; Moore, R. et al., 1987, J. Virol. 61:480-490). Evaluation of the clones indicated that homology to MMTV env gene varied from 95% to 99%. Another seven clones comprising only 250 bp were also sequenced.
30 Homology to MMTV env gene varied from 95% to 99% (data not shown). When compared to the human endogenous provirus HERV-K10, the homology of all the clones was less than 15%. When compared against all known viral and human genes (more than 130,000 entries) using the
35 lBl MacVector GenBank and GCG programs, the highest homology recorded was 18%.

6.2.4. Southern Blot Analysis Using Cloned Sequences

To investigate whether the env gene-like sequences were present in human DNA, Southern blot hybridization was performed using the cloned sequence as probe. DNAs from normal breast tissues, env positive or negative breast tumors, tumors other than breast and breast cancer cell lines were restricted with EcoRI and in some instances with PstI, BglII or KpnI. EcoRI is a frequent cutter restriction enzyme that digests MMTV proviral DNA between env and pol genes. Four different cloned 660 bp sequences were used as probes after labeling with ³²P by random prime-labeling. Results of some of the Southern blot hybridization experiments are shown in Fig. 5. They reveal the presence of a labeled restriction fragment migrating at approximately 7-8 kb in breast cancer DNA, in ED and two fragments in MCF-7 cells. Different restriction patterns were observed with the other three enzymes. The 660 bp sequence was absent in 10 normal tissues, 10 fibroadenomas and 10 tumors from other tissues. It is important to emphasize that hybridization conditions for these experiments were stringent (as described in Section 6.1) to avoid interference with endogenous sequences that might interact with the probes.

7. Example: Detection of a Retrovirus Proviral Fragment in Human Breast Cancer Cells and Tissues

7.1. Materials and Methods

To detect longer retrovirus proviral fragments in breast cancer samples, DNA was extracted from breast cancer carcinoma tissue samples as described above in Section 6.1. Two rounds of long PCR was performed on the DNA primers 5L (SEQ ID NO:11) and LTR3 (SEQ ID NO:19). The primer 5L contains nucleotides 7370-7395 of the MMTV BR6 genome (5'-3': CCAGATCGCCTTTAAGAAGG) (SEQ ID NO:11) and primer LTR3 contains nucleotides

9918-9927 of the MMTV BR6 genome (5'-3':
CGAACAGACACAAAGCGACG) (SEQ ID NO:19). Long PCR was
performed using protocols described by the manufacturer
(Perkin Elmer, Foster City, CA). The amplified
5 retroviral fragment isolated from the breast cancer
sample was cloned into the TA cloning vector
(Invitrogen) and automated sequencing was performed
as described in Section 6.1.

7.2 Results

10 An approximately 2.6 kb retroviral fragment
containing 1,228 bp of the 3' LTR sequence and 1,336 bp
of the env gene sequence of a potential provirus was
detected in a human breast carcinoma tissue sample by
the long PCR technique using the 5L and LTR3 primers.
15 The sequence of this retroviral fragment is shown in
Fig. 9. (SEQ ID NO:20).

When compared with the two strains of MMTV C3H
and BR6, the sequence homology was 90.8% and 90.7%,
respectively, over the MMTV genomic fragment from
20 nucleotides 7370-9937. When compared with the
endogenous retroviral sequences (HUMERKA), sequence
homology was only 58% in 36 bp and 71% in 74 bp.

8. Discussion

Search for virus-related sequences in human breast
25 cancer has been hampered by great variation reported
in previous studies, by the presence of endogenous
retroviral sequences in human DNA and by the lack of
sensitivity of the methods employed. The studies
reported herein circumvent these deficiencies by
30 focusing on sequences with low homology to human
endogenous retroviruses, by investigating a large
number of tumors and several types of controls and
by using the most sensitive technology presently
available.

35 The results indicate that unique MMTV env gene
sequences were present in 38.5% of the breast cancer

samples analyzed and 39.7% of archival samples of breast cancer and that these sequences were absent in normal tissues including lymphocytes from patients with positive breast cancer and in cancers other than breast. Normal breast tissue and fibroadenomas had a low frequency (1.8 to 6.9%) of positive results. When cloned and sequenced, the sequences were found to be highly homologous to MMTV env gene, but not to the endogenous retroviral sequences. Furthermore, experiments in which the cloned amplified sequences were used for hybridization with DNA from breast cancer or normal tissues revealed that homologous DNA was only present in breast cancer DNA. The results also indicate that a human breast carcinoma sample contained an about 2.6 kb MMTV-like fragment comprised of 1,336 bp of the env gene and 1,228 bp of the 3' LTR.

The detection of MMTV env gene sequences in two fibroadenomas out of 29 and in two normal breast tissue samples out of 107 samples is of uncertain significance. Although such results could potentially be artifactual, and thus may represent false positives, they may alternatively indicate the presence of histologically unrecognized cells that were or will be neoplastic.

Ninety percent (90%) of the breast cancers tested were invasive ductal carcinomas, which reflects the prevalence of this type of neoplasm. Most patients were node-positive which is probably artifactual since it was necessary that tumor size be sufficiently large to provide an aliquot for research and tumor size correlates with node positivity.

It is unlikely that differences in homology between MMTV env gene and the cloned human sequences are generated by errors committed by the Taq polymerase. It has been estimated that the rate of nucleotide misincorporation is 1×10^{-5} per cycle (Ehrlich et al, 1991, Science 252:1643-1651) and

therefore, only a total of 0.32 nucleotides
misincorporated should be expected in 660 bp after
50 cycles. The differences in homology between clones
from different patients is likely to represent
5 heterogeneity of the env gene.

In contrast to earlier, ambiguous data associating
MMTV-like sequences with human breast cancer, we have
clearly demonstrated the existence of such sequences in
breast cancer cells which cannot be explained by any
10 known human endogenous retroviral sequence. Our data do
not support the results of earlier studies which
indicated that, as in the mouse, MMTV-like sequences
were found in lymphocytes from two patients with breast
cancer (Crepin, M. et al., 1984, Biochem. Biophys. Res.
15 Comm. 118:324-331). The absence of MMTV env-like
sequences in lymphocytes could reflect the fate of a
unique lymphocyte subset over decades between initial
encounter and the appearance of clinical breast cancer;
alternatively, the human disease may differ from the
20 mouse model. Results from attempts to identify unique
MMTV-like pol gene sequences have shown that they
cannot be distinguished from the reverse transcriptase
sequences of endogenous retroviruses (Deen, K.C. and
Sweet, R.W., 1986, J. Virol. 57:422-432).

25 The origin of the MMTV env gene-like and 3' LTR-
like sequences found in tumor DNA could be the result
of integrated MMTV-like sequences from a human mammary
tumor virus. Polymorphism of endogenous retroviral
sequences is conceivable but can be ruled out because
30 these sequences were not detected in lymphocytes from
the positive patients, in sections of the cancerous
breast from which abnormal cells were absent, or in
normal breast tissue from patients with MMTV env-like
positive tumors. Recombination during tumorigenesis
35 between endogenous sequences to resemble the MMTV env
genes seems highly unlikely since no known gene or
viral sequence is more than 18% homologous to the

660 bp sequence. The longer about 2.6 kb MMTV-like fragment detected in a human breast carcinoma had minimal homology (58% in 36 bp and 71% in 74 bp) to endogenous human retroviral sequences. Thus, the most conservative interpretation is that our findings represent exogenous sequences from an agent similar to MMTV. Recombination between endogenous and exogenous env gene sequences are known to accelerate the development of malignancies in mice (DiFronzo, N.L. and Holland, C.A., 1993, J. Virol. 67:3763-3770). Whether the MMTV-like sequences belong to an entire acquired provirus or to an exogenous fragment integrated into endogenous sequences, is presently not known. Experiments are in progress to distinguish between these possibilities.

Several genetic alterations have been identified in human breast cancer that can be useful as markers for prevention, detection or prognosis (reviewed in Runnenbaum, I. et al., 1991, Proc. Natl. Acad. Sci. USA 88:10657-10661). The BRCA1 and BRCA2 genes have recently been described. They account for at least 5% of breast cancer and are related to familial breast cancer (Miki, Y. et al., 1994, Science 266:66-71; Wooster, R. et al., 1994, Science 265:2088-2090). We have primary evidence that familial clustering of the MMTV env gene-like sequences occurs, accounting for an even higher percentage of cancers in affected families (Holland et al. 1994, Proc. Am. Assoc. Cancer Res 35:218). The presence of MMTV-like sequences may be correlated with special clinical disease status, may provide another potential molecular marker, and may distinguish a subset of human breast cancer for which viral etiology is tenable. This has implications for epidemiology, therapy and prevention.

Various publications are cited herein, the contents of which are hereby incorporated by reference in their entireties.

SEQUENCE LISTING

(1) GENERAL INFORMATION

(i) APPLICANT: HOLLAND, JAMES

5 (ii) TITLE OF THE INVENTION: DETECTION OF MAMMARY TUMOR VIRUS-LIKE
SEQUENCES IN HUMAN BREAST CANCER

(iii) NUMBER OF SEQUENCES: 20

(iv) CORRESPONDENCE ADDRESS:

10 (A) ADDRESSEE: Brumbaugh, Graves, Donohue & Raymond
(B) STREET: 30 Rockefeller Plaza
(C) CITY: New York
(D) STATE: NY
(E) COUNTRY: USA
(F) ZIP: 10112-0228

15 (v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Diskette
(B) COMPUTER: IBM Compatible
(C) OPERATING SYSTEM: DOS
(D) SOFTWARE: FastSEQ Version 1.5

20 (vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER: NOT YET ASSIGNED
(B) FILING DATE: 08-NOV-1996
(C) CLASSIFICATION:

25 (vii) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER 08/555,394
(B) FILING DATE: 09-NOV-1995

(viii) ATTORNEY/AGENT INFORMATION:

30 (A) NAME: Kole, Lisa B
(B) REGISTRATION NUMBER: 35,225
(C) REFERENCE/DOCKET NUMBER: 30363-PCT - 165/

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: 212-408-2628
(B) TELEFAX: 212-765-2519
(C) TELEX:

35 (2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

40 (A) LENGTH: 15 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

45 (vi) ORIGINAL SOURCE:

(ix) FEATURE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

CCTCACTGCC AGATC

15

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 22 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(ix) FEATURE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

GGGAATTCCT CACTGCCAGA TC

22

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 19 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(ix) FEATURE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

CCTCACTGCC AGATCGCCT

19

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 18 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(ix) FEATURE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

TACATCTGCC TGTGTTAC

18

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

- 5 (ii) MOLECULE TYPE: cDNA
(iii) HYPOTHETICAL: NO
(iv) ANTISENSE: NO
(v) FRAGMENT TYPE:
(vi) ORIGINAL SOURCE:
(ix) FEATURE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

CCTACATCTG CCTGTGTTAC

20

(2) INFORMATION FOR SEQ ID NO:6:

- 10 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 15 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

- 15 (ii) MOLECULE TYPE: cDNA
(iii) HYPOTHETICAL: NO
(iv) ANTISENSE: NO
(v) FRAGMENT TYPE:
(vi) ORIGINAL SOURCE:
20 (ix) FEATURE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

CCGCCATACG TGCTG

15

(2) INFORMATION FOR SEQ ID NO:7:

- 25 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 15 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

- 30 (ii) MOLECULE TYPE: cDNA
(iii) HYPOTHETICAL: NO
(iv) ANTISENSE: NO
(v) FRAGMENT TYPE:
(vi) ORIGINAL SOURCE:
(ix) FEATURE:

35 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

ATCTGTGGCA TACCT

15

(2) INFORMATION FOR SEQ ID NO:8:

- 40 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 23 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

- 45 (ii) MOLECULE TYPE: cDNA
(iii) HYPOTHETICAL: NO
(iv) ANTISENSE: NO
(v) FRAGMENT TYPE:
(vi) ORIGINAL SOURCE:
(ix) FEATURE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

GGGAATTCAT CTGTGGCATA CCT

23

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(ix) FEATURE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

ATCTGTGGCA TACCTAAAGG

20

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 15 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(ix) FEATURE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

GAATCGCTTG GCTCG

15

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(ix) FEATURE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

CCAGATCGCC TTTAAGAAGG

20

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(ix) FEATURE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

TACAGGTAGC AGCACGTATG

20

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: N-terminal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

Leu Lys Arg Pro Gly Phe Gln Glu His Glu Met Ile
1 5 10

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE: N-terminal

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

Gly Leu Pro His Leu Ile Asp Ile Glu Lys Arg Gly
1 5 10

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: peptide
 (iii) HYPOTHETICAL: NO
 (iv) ANTISENSE: NO
 (v) FRAGMENT TYPE: N-terminal
 (vi) ORIGINAL SOURCE:
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

Thr Asn Cys Leu Asp Ser Ser Ala Tyr Asp Thr Ala
 1 5 10

(2) INFORMATION FOR SEQ ID NO:16:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 10 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: peptide
 (iii) HYPOTHETICAL: NO
 (iv) ANTISENSE: NO
 (v) FRAGMENT TYPE: N-terminal
 (vi) ORIGINAL SOURCE:

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

Asp Ile Gly Asp Glu Pro Trp Phe Asp Asp
 1 5 10

(2) INFORMATION FOR SEQ ID NO:17:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 662 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: cDNA
 (iii) HYPOTHETICAL: NO
 (iv) ANTISENSE: NO
 (v) FRAGMENT TYPE:
 (vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

| | | |
|----|--|-----|
| 35 | TCCTCACTGC CAGATCGCCT TTAAGAAGGA CGCCTTCTGG GAGGGAGACG AGTCTGCTCC | 60 |
| | TCCACGGTGG TTGCCTTGCG CCTTCCCTGA CCAAGGGGTG AGTTTTTCTC CAAAAGGGGC | 120 |
| | CCTTGGGTTA CTTTGGGATT TCTCCCTTCC CTCGCCTAGT GTAGATCAGT CAGATCAGAT | 180 |
| | TAAAAGCAAA AAGGATCTAT TTGGAAATTA TACTCCCCCA GTCAATAAAG AGGTTTCATCG | 240 |
| | ATGGTATGAA GCAGGATGGG TAGAACCTAC ATGGTTCTGG GAAAATTCTC CTAAGGATCC | 300 |
| 40 | CAATGATAGA GATTTTACTG CTCTAGTTCC CATAAGAAT TGTTTCGCTT AGTTGCAGCC | 360 |
| | TCAAGATATC TTATTCTCAA AAGGCAGGAT TTCAGGAACA TGAGATGATT CCTACATCTC | 420 |
| | TGTGTTACTT ACCCTTATGT CATATTATTA GGATTACCTC AGCTAATAGA TATAGAGAAA | 480 |
| | GAGGATCTAC TTTTCATATT TCCTGTTCTT CTTGTAGATT GACTAATTGT TTAGATTCTT | 540 |
| | CTGCCTACGA CTATGCAGCG ATCATAGTCA AGAGGCCGCC ATACGTGCTG CTACCTGTAG | 600 |
| 45 | ATATTGGTGA TGAACCATGG TTTGATGATT CTGCCATTCA AACCTTAGG TATGCCACAG | 660 |
| | AT | 662 |

(2) INFORMATION FOR SEQ ID NO:18:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 663 base pairs
 (B) TYPE: nucleic acid

(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA
(iii) HYPOTHETICAL: NO
(iv) ANTISENSE: NO
(v) FRAGMENT TYPE:
(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

| | | | | | | | |
|----|------------|------------|-------------|------------|------------|-------------|-----|
| 10 | TCCTCACTGN | CAGATCGCCT | TTAAGAAGGA | CGCCTTCTGG | GAGGGAGACG | AGTCTGCTCC | 60 |
| | TCCACGGTGG | TTGACTTGCG | CCTTCCCTGA | CCAGGGGGTG | AGTTTTTCTC | CAAAAGGGGC | 120 |
| | CCTTGGGTTA | CTTGGGATT | TCTCCCTTCC | CTCGCCTAGT | GTAGATCAGT | CAGATCAGAT | 180 |
| | TAAAGCAAA | AAGGATCTAT | TTGGAAATTA | TACTCCCCCT | GTCAATAAAG | AGGTTTCATCG | 240 |
| | ATGGTATGAA | GCAGGATGGG | TAGAACCCTAC | ATGGTTCTGG | GAAAATTCTC | CTAAGGATCC | 300 |
| | CAATGATAGA | GATTTTACTG | CTCTAGTTCC | CATACAGAAT | TGTTTCGCTT | AGTTGCAGCC | 360 |
| 15 | TCAAGATATC | TTATTCACAA | AAGGCAGGAT | TTCAAGAACA | TGACATGAAT | CCCTACATCT | 420 |
| | CTGTGTTACT | TACCCTTATG | CCANANTATT | AGGATTACCT | CAGCTAATAG | ATATAGAGGA | 480 |
| | AGAGGATCTA | CTTTTCATAT | TTCTGTTTCT | TCTGTAGAT | TGACTAATTG | TTTAGATTCT | 540 |
| | TCTGCCTACG | ACTATGCAGC | GATCATAGTC | AAGAGGCCGC | CATACGTGCT | GCTACCTGTA | 600 |
| 20 | GATATTGGTG | ATGAACCATG | GTTTGATGAN | NCTGCCANTC | AAACCTTTAG | GTATNCCACA | 660 |
| | GAT | | | | | | 663 |

(2) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA
(iii) HYPOTHETICAL: NO
(iv) ANTISENSE: NO
(v) FRAGMENT TYPE:
(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

CGAACAGACA CAAACACACG

20

(2) INFORMATION FOR SEQ ID NO:20:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 2598 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA
(iii) HYPOTHETICAL: NO
(iv) ANTISENSE: NO
(v) FRAGMENT TYPE:
(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

| | | | | | | | |
|----|-------------|-------------|------------|------------|------------|------------|------|
| | CGAACAGACA | CAAACACACG | AGAGGTGAAT | GTTAGGACTG | TTGCAAGTTT | ACTCAAAAAA | 60 |
| | CAGCACTCTT | TTATATCATG | GTTTACATAA | GCATTTACAT | AAGACTTGGA | TAAGTTCCAA | 120 |
| | AAGAACATAG | GAGAATAGAA | CACTCAGAGC | TTAGATCAAA | ACATTTGATA | CCAAACCAAG | 180 |
| 5 | TCAGGAAACC | ACTTGTCTCA | CATCCTTGTT | TTAAGAACAG | TTTGTGACCC | TGAACTTACT | 240 |
| | TAAACCTTGG | GAACCGCAAN | GTTGGGCTCA | TAAAGGTTAT | CCATTATAGC | TCATGCCAAA | 300 |
| | ATTATCTGCA | GAATGTGTT | CCTAATGTC | TAGCCACTGC | CCCCTCCCTT | GGTATAATGA | 360 |
| | AAATCCTTCC | CCCAACGTTT | ATCCCACTCC | CCTAGATAAA | TATAATCATG | TACCTGTTGT | 420 |
| | TTTATGTGCT | CTTTTCTTTC | CTGAGTTAAC | ACACACCAAG | GAGGTCTAGC | TCTGGCGAGT | 480 |
| 10 | CTTTCACGAA | AGGGGAGGGA | TCTGTACAAC | ACTTTATAGC | CGTTGACTGT | GACCCACCTA | 540 |
| | TCGAAATTTA | AATCGTATCT | TCCTGTATAT | GCTAGCGGGG | CGTCTGTTGG | TCTGTAGATG | 600 |
| | TAAGTCCCGG | TTGCCACCAC | TCGTCTCCTA | TTTTGACAAG | CGTACTCCTC | TTTCCCCTTT | 660 |
| | TTACTTCTAG | GCCTGAGGCC | CTTAGTCCTT | GCACCTGTTC | TTCAACTGAG | GTTGAGCGTC | 720 |
| | TCTTTCTATT | TTCTATTCCC | ATTTCTAACC | TTTGAATTTG | AGTAAATATA | GTGCTAAAAG | 780 |
| 15 | ACAAAGATTC | ATTTCTTAAC | ATCATGATTA | ATAATCGACC | TATTGGATTG | GTCTTATTGG | 840 |
| | TAAAAATATA | ATTTTTAGCA | AGCATTCTTA | TTTCTATTTC | TGAAGGACAA | AGTCGGTGTG | 900 |
| | GCTTGTAAANA | GGAANTTGGC | TGTGGTCTTT | GCCCCACGAG | GAAGGTCGAG | TTCTCCGAAT | 960 |
| | TGTTTTAGATT | GTAATCTTGC | ACAGAAGAGT | TATTAAAGA | ATCAAGGGTG | AGAGCCCTGC | 1020 |
| | GAGCACGAAC | CGCAACTTCC | CCCAATAGCC | CCAGGCAAAG | CAGAGCTATG | CCAAGTTTGC | 1080 |
| 20 | AGCAGANAAT | GAGTATGTCT | TTGTCTGATG | GGCTCATCCG | CGTGCACGCA | GACGGGTCGT | 1140 |
| | CCTTGGTGGG | AAACAACCCC | TTGGCTGCTT | CTCTCCTAAG | TGTAGGACAC | TCTCGGAGT | 1200 |
| | TCAACCATTT | CTGCTGCAGG | CGCGGCATTT | CCCCCTTTT | TCTTTTAA | AAGAAGCAGC | 1260 |
| | TTAAGATCTG | ACTGCACTTG | GTCAAGGCTC | TTCCGAAAGC | ACTGGAAAT | AACGGGAAA | 1320 |
| | ATCATAGTA | CTATGACCAA | AAGCAGGCTC | CCAACCTCTA | TAAAAATGAA | ATATTGTGTT | 1380 |
| 25 | CTAATCCAAT | GGATTTAAAG | CCTTTACTCC | ATTGGCNAAG | GANTGANCCA | ACCCCTGAGG | 1440 |
| | TCCCTGCGTT | CAAATTTTTT | TGCTCNTATC | CTAATCCAAT | TGGTAACCCC | GTTTNTTTTT | 1500 |
| | GAAACTCATG | TCTTCAAATG | CCCAATAAAT | GAGCCCTGGT | TCTTTCCCAG | CTCTCAGAAG | 1560 |
| | CATTATACGG | NANAGGTGTG | ACACAGCATA | AAATCATAAT | TTGCATGACA | CCTAGTGGAC | 1620 |
| | ATTCTGGTCT | TTAAGTTTGC | CACATCTTGT | CCCAACTCTA | AAACTACTTC | TTCTAAAGCA | 1680 |
| 30 | TTAAGTCTAG | CTTTCAATTT | TAAGTCTATT | ATTCTTTGTT | CAGATNAGGC | TAATGTAACA | 1740 |
| | TTTCTATGAA | GATTATTAAC | AAACGTAGCA | GTTTGCATCT | CCTTAACTAA | GGCAGTAGTA | 1800 |
| | GCTACAGCAA | AGGAAGTGAT | AATAGCAATT | AAAGCAGATA | TGCCCAGAAT | AATGGCAGCG | 1860 |
| | ACGAATCGCT | TAGCTCGAAT | TAAATCTGTG | GCATACCTAA | AGGTTTGAAT | GGCAGAATCA | 1920 |
| | TCAAACCATG | GTTTCATCACC | AATATCTACA | GGTTACAACA | CATATGGCGG | CCCCTTGAAT | 1980 |
| 35 | ATGAATCGCT | GCATATCCGT | NGGCAAAAAA | TCTAACCATT | ATTCCTCCTN | CCNAAAAACG | 2040 |
| | GGATTTGAAA | NTTATNCCC | TTNCCCCNAA | CCCANACCGA | GGTACCCCAT | AATNGGGGGG | 2100 |
| | GTATCTANAA | NAGGGCATAG | GGGTAAGAAA | AACGGCAGAG | NGGGATCNTT | TATGTTCNNG | 2160 |
| | AAATTCNNGG | TTTGGGAGAA | TAAGATTCTG | GAGGCTGCAA | ATTAAGGGAA | ACATTNTGTA | 2220 |
| | TGGGGAATAG | AGCAGTAAAA | TCTCTATCAT | GGGGATCTTT | AGGGAGAATT | TTCCCAGGAA | 2280 |
| 40 | CCAAGTAGGT | TCNAACCCAT | CNTGCTTCAT | ACCATCGATG | AACNTCTTTA | TTGACAGGGG | 2340 |
| | GAGTATAATT | TCCAAATAGA | TCCTTTTGT | TTTTAATCTG | ATCTGACTGA | TCTACACTAG | 2400 |
| | GCGGGGAAG | GGAGAAATCC | CAAAGTAACC | CAAGGGCCCC | TTTTGGAGAA | AAACTCACCC | 2460 |
| | CCTGGTCAGG | GAAGGCGCAA | GGCAACCACC | GTGGAGGAGC | AGACTCGTCT | CCCTCCCGA | 2520 |
| 45 | AGGCGTCCCT | CTTAAAGGCG | ATCTGGAGGA | GCAGACTCGT | CTCCCTCCCA | GAAGGCGTCC | 2580 |
| | TTCTTAAAGG | CGATCTGG | | | | | 2598 |

Claims

- 1 1. A composition comprising an oligonucleotide primer
2 which may be used to detect the presence of a
3 nucleic acid molecule which (i) hybridizes to the
4 env gene of a mouse mammary tumor virus; (ii) is
5 present in at least 38 percent of DNA samples
6 prepared from breast cancer tissue of different
7 human subjects; and (iii) hybridizes to less than
8 7 percent of DNA samples prepared from tissues
9 other than breast cancer tissue from different
10 human subjects.
- 1 2. The composition of claim 1, wherein the
2 oligonucleotide primer comprises the sequence
3 CCTCACTGCCAGATC (SEQ ID NO:1).
- 1 3. The composition of claim 1, wherein the
2 oligonucleotide primer comprises the sequence
3 GGAATTCCTCACTGCCAGATC (SEQ ID NO:2).
- 1 4. The composition of claim 1, wherein the
2 oligonucleotide primer comprises the sequence
3 CCTCACTGCCAGATCGCCT (SEQ ID NO:3).
- 1 5. The composition of claim 1, wherein the
2 oligonucleotide primer comprises the sequence
3 TACATCTGCCTGTGTTAC (SEQ ID NO:4).
- 1 6. The composition of claim 1, wherein the
2 oligonucleotide primer comprises the sequence
3 CCTACATCTGCCTGTGTTAC (SEQ ID NO:5).
- 1 7. The composition of claim 1, wherein the
2 oligonucleotide primer comprises the sequence
3 CCGCCATACGTGCTG (SEQ ID NO:6).

- 1 8. The composition of claim 1, wherein the
2 oligonucleotide primer comprises the sequence
3 ATCTGTGGCATACT (SEQ ID NO:7).
- 1 9. The composition of claim 1, wherein the
2 oligonucleotide primer comprises the sequence
3 GGGAATTCATCTGTGGCATACT (SEQ ID NO:8).
- 1 10. The composition of claim 1, wherein the
2 oligonucleotide primer comprises a sequence
3 selected from the group consisting of
4 ATCTGTGGCATACTAAAGG (SEQ ID NO:9);
5 GAATCGCTTGGCTCG (SEQ ID NO:10);
6 CCAGATCGCCTTTAAGAAGG (SEQ ID NO:11); and
7 TACAGGTAGCAGCACGTATG (SEQ ID NO:12).
- 1 11. An essentially purified peptide encoded by a
2 nucleic acid molecule which (i) hybridizes to
3 a gene of MMTV; (ii) is present in at least
4 20 percent of DNA samples prepared from breast
5 cancer tissue of different human subjects; and
6 (iii) is present in less than 5 percent of DNA
7 samples prepared from tissues other than breast
8 cancer tissue from different human subjects.
- 1 12. An antibody which specifically binds to the
2 peptide of claim 11.
- 1 13. The peptide according to claim 11 which comprises
2 the amino acid sequence LKRPGFQEHMI (SEQ ID
3 NO:13).
- 1 14. An antibody which specifically binds to the
2 peptide of claim 13.
- 1 15. The peptide according to claim 11 which comprises
2 the amino acid sequence GLPHLIDIEKRG (SEQ ID NO:14).

1 16. A method of diagnosing breast cancer in a human
2 subject, comprising detecting the presence of a
3 peptide encoded by a nucleic acid molecule which
4 (i) hybridizes to the env gene of 3' LTR of a
5 mouse mammary tumor virus; (ii) is present in at
6 least 20 percent of DNA samples prepared from
7 breast cancer tissue of different human subjects;
8 and (iii) is present in less than 5 percent of DNA
9 samples prepared from tissues other than breast
10 cancer tissue from different human subjects.

1 17. The method according to claim 16, wherein the
2 presence of a peptide comprising the amino acid
3 sequence LKRPGFQEHMI (SEQ ID NO:13) is detected
4 by the binding of an antibody specific to the
5 peptide.

1 18. The method according to claim 16, wherein the
2 presence of a peptide comprising the amino acid
3 sequence GLPHLIDIEKRG (SEQ ID NO:14) is detected
4 by the binding of an antibody specific to the
5 peptide.

1 19. The method according to claim 16, wherein the
2 presence of a peptide comprising the amino acid
3 sequence TNCLDSSAYDTA (SEQ ID NO:15) is detected
4 by the binding of an antibody specific to the
5 peptide.

1 20. The method according to claim 16, wherein the
2 presence of a peptide comprising the amino acid
3 sequence DIGDEPWFD (SEQ ID NO:16) is detected by
4 the binding of an antibody specific to the
5 peptide.

1 21. A composition comprising an oligonucleotide primer
2 which may be used to detect the presence of a

3 nucleic acid molecule which (i) hybridizes to a
4 nucleic acid comprised of a sequence selected from
5 the group consisting of the env gene and the 3'
6 LTR of a mouse mammary tumor virus; (ii) is
7 present in a substantial percentage of DNA samples
8 prepared from breast cancer tissue of different
9 human subjects; and (iii) hybridizes to less than
10 5 percent of DNA samples prepared from tissues
11 other than breast cancer tissue from different
12 human subjects.

1 22. The composition of claim 1, wherein the
2 oligonucleotide primer comprises the sequence
3 CCAGATCGCCTTTAAGAAGG (SEQ ID NO:11).

1 23. The composition of claim 1, wherein the
2 oligonucleotide primer comprises the sequence
3 CGAACAGACACAAACACACG (SEQ ID NO:19).




FIGURE 1




FIGURE 2




FIGURE 3





FIGURE 4




FIGURE 5

A B C D E F G H



FIGURE 6




FIGURE 7

NTTENV 980 | 1000 | 1020 |
[1810] TCCTCACTGACAGATGGCT TTAAGAAGGAGGCTTCTTC GAGGAGACAGATCTCTCTC
MS1627.Seq TCCTCACTGACAGATGGCT TTAAGAAGGAGGCTTCTTC GAGGAGACAGATCTCTCTC
5 10 15 20 25 30 35 40 45 50 55 60

NTTENV 1040 | 1060 | 1080 |
[1810] TCACGGTGGTTGACTTGG CCTTCCTGACCAAGGGCTG AGTTTCTCCAAAAGGGCC
MS1627.Seq TCACGGTGGTTGACTTGG CCTTCCTGACCAAGGGCTG AGTTTCTCCAAAAGGGCC
65 70 75 80 85 90 95 100 105 110 115 120

NTTENV 1100 | 1120 | 1140 |
[1810] CCTTGGGTTACTTTGGGAT TCTCCCTTCCCTGGCTAGT GTAGATCACTCAGATCAGAT
MS1627.Seq CCTTGGGTTACTTTGGGAT TCTCCCTTCCCTGGCTAGT GTAGATCACTCAGATCAGAT
125 130 135 140 145 150 155 160 165 170 175 180

NTTENV 1160 | 1180 | 1200 |
[1810] TAAAGCAAAAAGATCTAT TTGGAATTATCTCCCGCA GTCAATAAGAGGTTTCATC
MS1627.Seq TAAAGCAAAAAGATCTAT TTGGAATTATCTCCCGCT GTCAATAAGAGGTTTCATC
185 190 195 200 205 210 215 220 225 230 235 240

NTTENV 1220 | 1240 | 1260 |
[1810] ATGGTATGAAGCAGGATGG TGAAGCTACATGGTTCTGG GAAATTCCTTAAGGATCC
MS1627.Seq ATGGTATGAAGCAGGATGG TGAAGCTACATGGTTCTGG GAAATTCCTTAAGGATCC
245 250 255 260 265 270 275 280 285 290 295 300

NTTENV 1280 | 1300 | 1320 |
[1810] CAATGATAGAGATTTTACTG CTCTAGTTCCCATACAGAT TGTTCCTCTAGTTCCAGCC
MS1627.Seq CAATGATAGAGATTTTACTG CTCTAGTTCCCATACAGAT TGTTCCTCTAGTTCCAGCC
305 310 315 320 325 330 335 340 345 350 355 360

NTTENV 1340 | 1360 | 1380 |
[1810] TCAAGATATCTTATTCACAA AAGGCAGGATTTCAAGACA TCAGATG-ATCCCTACATCT
MS1627.Seq TCAAGATATCTTATTCACAA AAGGCAGGATTTCAAGACA TCAGATGAATCCCTACATCT
365 370 375 380 385 390 395 400 405 410 415 420

NTTENV 1400 | 1420 | 1440 |
[1810] CTGTGTACTTACCTTTATC CCAGCTATTAAGATTAAGT CAGCTATAGATATAGAAA
MS1627.Seq CTGTGTACTTACCTTTATC CCAGCTATTAAGATTAAGT CAGCTATAGATATAGAAA
425 430 435 440 445 450 455 460 465 470 475 480

NTTENV 1460 | 1480 | 1500 |
[1810] AGAGCATCTACTTTTCAAT TTCTGTCTCTCTTTAGAT TGACTAATTTTATGATCT
MS1627.Seq AGAGCATCTACTTTTCAAT TTCTGTCTCTCTTTAGAT TGACTAATTTTATGATCT
485 490 495 500 505 510 515 520 525 530 535 540

NTTENV 1520 | 1540 | 1560 |
[1810] TCTGCTAGACTATGCAGC GATCATAGTCAGAGGCGCC CATACCTCTCTACTCTTA
MS1627.Seq TCTGCTAGACTATGCAGC GATCATAGTCAGAGGCGCC CATACCTCTCTACTCTTA
545 550 555 560 565 570 575 580 585 590 595 600

NTTENV 1580 | 1600 | 1620 | 1640
[1810] GATATGTTGATGACCAATG GTTGTATGATCTGCCATTC AAACCTTAGCTATGCCA GAT
MS1627.Seq GATATGTTGATGACCAATG GTTGTATGATCTGCCATTC AAACCTTAGCTATGCCA GAT
605 610 615 620 625 630 635 640 645 650 655 660

FIGURE 8

CGAACAGACACACACACACGAGAGGTGAATGTTAGGACCTTGCAAGTTTA
CTCAAAAAACAGCACTCTTTTATATCATGGTTTACATAAGCATTACATAAGA
CTTGGATAAGTTCCAAAAGAACATAGGAGAATAGAACACTCAGAGCTTAGAT
CAAAACATTTGATACCAAACCAAGTCAGGAAACCACTTGTCTCACATCCTTG
TTTTAAGAACAGTTTGTGACCCTGAACCTTAAACCTTGGGAACCGCAAN
GTTGGGCTCATAAAGGTTATCCATTATAGCTCATGCCAAAATTATCTGCAGA
AATGTGTTCCCTAATTGTCTAGCCACTGCCCTCCCTTGGTATAATGAAAAT
CTTTCCCCAACGTTTCATCCCACTCCCTAGATAAATAATCATGTACCTGT
TGTTTTATGTCGTCTTTTTCTTCCTGAGTTAACACACACCAAGGAGGTCTAGC
TCTGGCGAGTCTTTCACGAAAGGGGAGGGATCTGTACAACACTTTATAGCC
GTTGACTGTGACCCACCTATCGAAATTTAAATCGTATCTTCCTGTATATGGTA
GCGGGGCGTCTGTTGGTCTGTAGATGTAAGTCCCGGTTGCCACCACCTGTC
TCCTATTTTGACAAGCGTACTCCTCTTTCCCCTTTTACTTCTAGGCCTGAGG
CCCTTAGTCCTTGACCTGTTCTTCAACTGAGGTTGAGCGTCTCTTTCTATTT
TCTATTCCCATTCTAACCTTTGAATTTGAGTAAATATAGTGCTAAAAGACAA
AGATTCAATTTCTTAACATCATGATTAATAATCGACCTATTGGATTGCTTTATT
GGTAAAAATATAATTTTAGCAAGCATTCTTATTTCTATTTCTGAAGGACAAA
GTCCGGTGTGGCTTGTAANAGGAANTTGGCTGTGGTCCTTGCCCCACGAGGA
AGGTTCGAGTTCTCCGAATTGTTTAGATTGTAATCTTGACAGAAAGAGTTATTA
AAAGAATCAAGGGTGAGAGCCCTGCGAGCACGAACCGCAACTTCCCCCAAT
AGCCCCAGGCAAAGCAGAGCTATGCCAAGTTTGCAGCAGANAATGAGTATG
TCTTTGTCTGATGGGCTCATCCGCGTGACGCGAGACGGGTCTCCTTGGTG
GGAAACAACCCCTTGGCTGCTTCTCTCCTAAGTGTAAGGACACTCTCGGGAG
TTCAACCATTCTGCTGCAGGCGCGGCATTTCCCCTTTTTCTTTTTTAAAA
GAAGCACGTTAAGATCTGACTGCACCTGGTCAAGGCTCTTCGCAAGCACT
GGAAAATAACGGGGAAAATCATAAGTACTATGACCAAAGCAGGGCTCCAA
CTCCTATAAAAATGAAATATTGTGTTCTAATCCAATGGATTTAAAGCCTTAC
TCCATTGGCNAAGGANTGANCCAACCCCTGAGGTCCCTGCGTTCAATTTTT
TTGCTCNTATCCTAATCCAATTGGTAACCCCGTTTNTTTTTGAAACTCATGTC
TTCAAATGCCAATAAATGAGCCCTGGTTCTTTCCCAGCTCTCAGAAGCATT
ATACGGNANAGGTGTGACACAGCATAAAATCATAATTTGCATGACACCTAGT
GGACATTCTGGTCTTTAAGTTTGCCACATCTTGTCCCAACTCTAAACTACTT
CTTCTAAAGCATTAAAGTCTAGCTTTCAATTTAAGTCTATTATTCTTTGTTTCA
ATNAGGCTAATGTAACATTTCTATGAAGATTATTAACAAACGTAGCAGTTTGC
ATCTCCTTAACCTAAGGCAGTAGTAGCTACAGCAAAGGAAGTGATAATAGCAA
TTAAAGCAGATATGCCCAGAATAATGGCAGCGACGAATCGCTTAGCTCGAAT
TAAATCTGTGGCATAACCTAAAGGTTTGAATGGCAGAATCATCAAAACCATGGT
TCATCACCAATATCTACAGGTTACAACACATATGGCGGCCCTTGAATATGA
ATCGCTGCATATCCGTNGGCAAAAAATCTAACCATTATTCCTCCTNCCNAAA
AACGGGATTTGAAANTTATNCCCCTTNCCCCNAACCCANACCGAGGTACCC
CATAATGNNGGGGGGTATCTANAANAGGGCATAGGGGTAAGAAAAACGGCA
GAGNNGGATCNTTTATGTTTNGGAAATTCNGGGTTTGGGAGAATAAGATTCT
GGAGGCTGCAAAATTAAGGGAAACATTNTGTATGGGGAATAGAGCAGTAAAA
TCTCTATCATGGGGATCTTTAGGGAGAATTTTCCAGGAACCAAGTAGGTTCT
NAACCCATCNTGCTTCATACCATCGATGAACNTCTTTATTGACAGGGGGAGT
ATAATTTCCAAATAGATCCTTTTGTTTTAAATCTGATCTGACTGATCTACACT
AGGCGGGGGAAGGGAGAAATCCCAAAGTAACCCAAGGGCCCCCTTTTGGAG
AAAAACTCACCCCTGGTCAGGGAAGGCGCAAGGCAACCACCGTGGAGGA
GCAGACTCGTCTCCCTCCCAGAAGGCGTCCTTCTTAAAGGCGATCTGGAGG
AGCAGACTCGTCTCCCTCCCAGAAGGCGTCCTTCTTAAAGGCGATCTGG

FIGURE 9

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US96/17877

| A. CLASSIFICATION OF SUBJECT MATTER IPC(6) : Please See Extra Sheet. US CL : 435/6, 5, 91.2, 7.1, 7.2; 536 23.1, 24.3, 24.33; 530/388.1, 300 According to International Patent Classification (IPC) or to both national classification and IPC | | |
|--|---|--|
| B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) U.S. : 435/6, 5, 91.2, 7.1, 7.2; 536 23.1, 24.3, 24.33; 530/388.1, 300 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) Please See Extra Sheet. | | |
| C. DOCUMENTS CONSIDERED TO BE RELEVANT | | |
| Category* | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
| A | REDMOND et al. Sequence and expression of the mouse mammary tumour virus env gene. The EMBO Journal. 1983, Volume 2, Number 1, pages 125-131. See entire document. | 1-20 |
| A | FAFF et al., Retrovirus-like particles from the human T47D cell lines are related to mouse mammary tumour virus and are of human endogenous origin. Journal of General Virology. 21 May 1992, Volume 73, pages 1087-1097. See abstract. | 1-20 |
| <input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex. | | |
| * Special categories of cited documents: | *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention *A* document defining the general state of the art which is not considered to be of particular relevance *E* earlier document published on or after the international filing date *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) *O* document referring to an oral disclosure, use, exhibition or other means *P* document published prior to the international filing date but later than the priority date claimed *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art *Z* document member of the same patent family | |
| Date of the actual completion of the international search 04 FEBRUARY 1997 | | Date of mailing of the international search report 18 MAR 1997 |
| Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230 | | Authorized officer DIANNE REES <i>JAB for</i> Telephone No. (703) 308-0196 |

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US96/17877

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

| Category* | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|-----------|--|-----------------------|
| A | CREPIN et al. Sequences Related to Mouse Mammary Tumor Virus Genome in Tumor Cells and Lymphocytes from Patients with Breast Cancer. Biochemical and Biophysical Research Communications. 13 January 1984, Volume 118, Number 1, pages 324-331. See entire document. | 1-20 |
| A | MESA-TEJADA et al. Detection in human breast carcinomas of an antigen immunologically related to a group-specific antigen of mouse mammary tumor virus. Proceedings of the National Academy of Sciences, USA. March 1978, Volume 75, Number 3, pages 1529-1533. | 1-20* |

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US96/17877

A. CLASSIFICATION OF SUBJECT MATTER:

IPC (6):

C12Q 1/68, 1/70; C12P 19/34; C07H 21/02, 21/04; G01N 33/53; C07K 15/28; 5/00

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

APS, BIOSIS, BIOTECHABS, BIOTECHDS, CABA, CAPLUS, CANCERLIT, DGENE, DRUGU, EMBASES, MEDLINE, USPATFULL, TOXLIT, TOXLINE, JAPIO, WPIDS
search terms: MMTV, mouse mammary tumor virus, PCR, hybridization, antibodies, immunoassays, Westerns, searched SEQ. ID. Nos.